

# Vorticity dynamics, Drift Wave Turbulence, and Zonal Flows: A Look Back and A Look Ahead

P.H. Diamond<sup>1,2</sup>

A. Hasegawa<sup>3</sup>

K. Mima<sup>4</sup>

<sup>1</sup> WCI Center for Fusion Theory, NFRI, Korea

<sup>2</sup> CMTFO and CASS, UCSD, USA

<sup>3</sup> Osaka University, Japan

<sup>4</sup> The Graduate School for the Creation of New Photonics Industries and ILE, Osaka University, Japan

**Abstract.** This paper surveys the basic ideas and results on fundamental models of drift wave turbulence, the formation of zonal flows, shear suppression of turbulence and transport, coupled drift wave and zonal flow dynamics, and application to transport bifurcations and transitions. Application to vortex dynamics and zonal flow phenomena in EMHD systems are discussed, as well. These are relevant to aspects of ICF and laser plasma physics. Throughout, an effort is made to focus on fundamental physics ideas.

## 1. Introduction

Inhomogeneity of plasma profiles in magnetically confined plasmas is responsible for generation of various instabilities that may lead to turbulent plasma states. Study of heat and particle confinement of plasmas in the presence of such turbulence is a fundamental issue for the success of controlled thermonuclear fusion. We address this issue in this manuscript. In the first part, we present the historical developments of the theory of strong electrostatic turbulence of magnetically confined plasmas and of the generation of azimuthal zonal flows. The model equation (the Hasegawa-Mima equation[5]) that includes plasma density inhomogeneity, nonlinear convection of plasma vortices, and Boltzmann electrons has two quadratic independent globally conserved quantities in the inviscid limit, energy and potential enstrophy. This fact indicates inverse cascade of turbulent energy spectra and the condensation of the spectrum to form a zonal flow in the azimuthal direction[10, 11]. Supporting simulation results are discussed.

The second part of the paper reviews the essential elements of the theory of drift wave-zonal flow turbulence[9]. The dynamics of zonal flow generation by potential vorticity or polarization charge transport[28] and the shearing feedback of zonal flows on the underlying microturbulence are discussed. We show that potential vorticity transport or mixing is fundamental to zonal flow formation. Shear decorrelation[30] physics is elucidated. Special attention is focused on the self-consistent feedback loop system - i.e. the predator-prey model[35, 37] - formed by the turbulence and flow shears. Simulation results[36] supporting the predator-prey model are reviewed. The relevance of these ideas to the ongoing quest to understand the  $L \rightarrow H$  transition and multi-predator prey models[37] relevant to the  $L \rightarrow H$  transition are discussed.

The third part deals with application to electron magnetohydro (EMHD) dynamics. EMHD[64] plays important roles in laser plasmas[65, 66, 67, 68, 69, 70], magnetic field reconnection plasmas[71, 72, 73, 74, 75, 76, 77] and space and astronomical plasma phenomena[78, 79] when the space and time scales are much shorter than the ion scales. The electro-magnetic fluctuations with frequencies much lower than the electron plasma frequency have structures like whistler waves. The turbulence of such electro-magnetic fluctuations are so called whistler turbulence. Examples of whistler turbulence are the turbulence predicted by the models of gamma ray burst[78, 79] and by the models for relativistic electron beam transport in plasmas[68, 69]. The whistler turbulence also plays important roles in collisionless magnetic reconnection in solar flare and magnetosphere[71, 72, 73, 74, 75, 76, 77].

The remainder of the paper is organized as follows. Section II deals with the formulation of the basic models and the basic aspects of zonal flow formation. Section III continues the discussion of zonal flow physics and formation, addresses shear suppression and describes the coupled dynamics of drift waves and zonal flows. Section IV presents application to EMHD vortices and structures. It also addresses applications of the ideas to problems in ICF and laser plasma physics. The last part of the paper is a brief

conclusion and outlook.

## 2. Drift Wave Vortex Models and Zonal Flows

### 2.1. Background

The study of plasma turbulence started almost immediately following the development of plasma physics, since plasma was recognized as intrinsically nonlinear. In the early stages, the weak turbulence theory based on perturbation analysis was popular, but starting in the late 1970's, when diagnosis based on electromagnetic wave scattering revealed strong turbulence property, attention to the theory of strong plasma turbulence increased. Hasegawa and Mima derived a model equation that describes the strong electrostatic turbulence based on the nonlinear convection of ion vortices that accompanies plasma density fluctuations. The equation demonstrates an existence of conservation of (potential) enstrophy (squared vorticity) in addition to energy. Analogy of this property to the two dimensional Navier-Stokes equation led the authors to predict an inverse spectral cascade and formation of self-organization of the turbulence. In the 1980's, Hasegawa together with Wakatani, extended the Hasegawa-Mima equation to include driving terms of the turbulence based on resistivity and magnetic field curvature and demonstrated the inverse cascade that leads to formation of the azimuthal zonal flow. They further predicted that the zonal flow could inhibit turbulent radial loss of the plasmas. Anomalous plasma loss due to turbulence has been the major concern in plasma confinement with a magnetic field, however, the prediction by Hasegawa and Wakatani proved a hope that plasma can be confined by the its own turbulent property. In this chapter, we also add some recent simulation results that support such possibilities.

### 2.2. Derivation of the Hasegawa-Mima Equation

When Mima was visiting Hasegawa at Bell Laboratories in 1976, we were introduced to very interesting laser scattering data by Slusher and Surko (1976)[1] obtained from density fluctuations of drift wave frequency range of a Princeton Tokamak. The data, in agreement with those obtained by Mazzucato (1976)[2] by microwave scattering, showed a very broad frequency spectrum whose width is wider than the expected drift frequency itself. Hasegawa and Mima were intrigued by the data and immediately constructed theory of drift wave turbulence that may account for this unexpected data. In response to answer criticism that the broad spectra might have been due the path averaging of the scattering date, Slusher and Surko (1978)[3] later have performed pin point scattering experiment by means of two lasers and demonstrated the validity of the earlier date.

We recognized that in order to account for the density fluctuation observed by those data, one needs to take into account of compressible ion flow. Since  $E \times B$  drift is incompressible, we recognized importance of the polarization drift, which is intrinsically nonlinear because of the convective electric field. We then realized that this nonlinear

polarization drift could best be recognized as nonlinear. The ion vorticity  $\Omega$  due to the  $E \times B$  drift can be describing of the Laplacian of the electrostatic potential  $\phi$ ,

$$\Omega = (\nabla \times \mathbf{v}_\perp) \cdot \hat{z} = \nabla \times \left( \frac{-\nabla\phi \times \hat{z}}{B_0} \right) \cdot \hat{z} = \frac{\nabla_\perp^2 \phi}{B_0} \quad (1)$$

Here,  $B_0$  is the flux density of the ambient magnetic field and  $\nabla_\perp^2$  is the Laplacian operator in the direction perpendicular to the magnetic field. The equation of ion vorticity can be constructed by taking a curl of the ion equation of motion in the direction normal to the ambient magnetic field. If the pressure and density gradients are parallel, the equation reads,

$$\frac{d}{dt}(\Omega + \omega_{ci}) + (\Omega + \omega_{ci})\nabla \cdot \mathbf{v}_\perp = 0 \quad (2)$$

Here  $\omega_{ci}$  is the ion cyclotron frequency. The compressible ion flow is related to the ion density fluctuation,  $n$ , through the continuity equation,

$$\nabla \cdot \mathbf{v}_\perp = -\frac{d}{dt} \ln n = -\frac{d}{dt} \left( \ln n_0 + \frac{e\phi}{T_e} \right) \quad (3)$$

Here, quasi-neutrality condition relates the ion density fluctuation to that of electrons, which will obey the Boltzmann equilibrium in this frequency range due to their inertia-less motions along the magnetic field. At this point the reader should recognize that even if we treat ions in two-dimensional plane normal to the direction of the magnetic field, we allow electron dynamics in three dimensions here.

From Eqns (2) and (3), we can construct the equation for ion vorticity for its compressible flow,

$$\frac{d}{dt} \left( \ln \frac{\omega_{ci}}{n_0} + \frac{\Omega}{\omega_{ci}} - \frac{e\phi}{T_e} \right) = \left( \frac{\partial}{\partial t} - \frac{\nabla\phi \times \hat{z}}{B_0} \cdot \nabla \right) \left( \ln \frac{\omega_{ci}}{n_0} + \frac{\Omega}{\omega_{ci}} - \frac{e\phi}{T_e} \right) \quad (4)$$

Equation (4) shows that the time evolution of vorticity and potential becomes nonlinear due to the nonlinear convection of the ion vorticity, as is expected, while the nonlinear convection of the potential does not contribute to nonlinear responses. Furthermore, if the spatial scale is on the order of the ion gyro radius at the electron temperature, which is the case of drift wave type fluctuations, the space time evolution of the potential field becomes fully nonlinear if the normalized potential field,  $e\phi/T_e$  becomes of the order of the ratio of drift wave frequency to the ion cyclotron frequency, which is very small.

Thus, for these space-time scales, the evolution of the potential field becomes fully nonlinear and is given by,

$$\frac{\partial}{\partial t} (\nabla_\perp^2 \phi - \phi) - \nabla\phi \times \hat{z} \cdot \nabla \left[ \nabla_\perp^2 \phi - \frac{1}{\epsilon} \ln \left( \frac{n_0}{\omega_{ci}} \right) \right] = 0 \quad (5)$$

Here the small parameter  $\epsilon$  represents the normalized amplitude as well as the time as shown below,

$$\epsilon \omega_{ci} t \equiv t, \quad x/\rho_s \equiv x, \quad e\phi/T_e \equiv \epsilon\phi, \quad \text{where } \rho_s = \sqrt{T_e/m_i}/\omega_{ci} \quad (6)$$

Equation (5) describes a fundamental property of electrostatic plasma turbulence in the drift wave frequency range in the absence of dissipations and often referred to as the Hasegawa-Mima equation[5].

### 2.3. Conservation laws, inverse cascades in kinetic and hydrodynamic regime

The model equation (5) that describes the drift wave turbulence demonstrates that drift wave becomes fully nonlinear even at a very low fluctuation level. In this section, we describe general properties of drift wave turbulence in hydrodynamic as well as in kinetic regimes based on this model equation.

*2.3.1. Conservation laws* Equation (5) has interesting conservation laws and its consequences. First it has energy conservation, which can be constructed by multiplying it with  $\phi$  and integrating over the volume,

$$\frac{\partial W}{\partial t} \equiv \frac{\partial}{\partial t} \int [(\nabla_{\perp} \phi)^2 + \phi^2] dV = 0 \quad (7)$$

Thus the sum of the kinetic and potential energies is conserved. The interesting part of the conservation law is that there exists additional conservation. The secondary conservation is the sum of the enstrophy, the squared vorticity, and the kinetic energy, sometimes called as potential enstrophy,

$$\frac{\partial U}{\partial t} \equiv \frac{\partial}{\partial t} \int [(\nabla_{\perp} \phi)^2 + (\nabla_{\perp}^2 \phi)^2] dV = 0 \quad (8)$$

The presence of conservation of the enstrophy in addition to the energy has been familiar in two-dimensional hydrodynamics and the interesting nature of its turbulence has been studied.

*2.3.2. Boltzmann statistics* In the absence of dissipation, the Boltzmann statistics that is obtained by minimizing the entropy in such a system has a strange character because of the additional constraint that comes from enstrophy conservation: the distribution function  $f$  that maximizes entropy with constraints of conservation of energy and enstrophy should read,

$$\delta \left( \int f \ln f dV - \lambda_1 W - \lambda_2 U \right) = 0 \quad (9)$$

Here  $\lambda$ s are Lagrange multipliers, and the resultant distribution function is given by

$$f \sim \exp(-\lambda_1 W - \lambda_2 U) \quad (10)$$

In the wave number space, since  $U_{\mathbf{k}} = k^2 W_{\mathbf{k}}$ , the average energy per mode becomes,

$$\langle W_{\mathbf{k}} \rangle = \frac{1}{\lambda_1 + k^2 \lambda_2} \quad (11)$$

As pointed out by Onsager (1949)[6], this result not only violates the equipartition law but also could results in a negative temperature if the product of  $\lambda_1$  and  $\lambda_2$  is negative, thus modal statistics can be strange, indicating consequences that may lead unexpected states.

*2.3.3. Self-organized state in hydrodynamic regime* In the presence of dissipation, turbulence described by the Hasegawa-Mima equation may lead to a dual cascade of the spectrum as pointed out by Kraichnan for two-dimensional Navier-Stokes turbulence. The inertial range Kolmogorov spectra (1949)[7] can be defined dually, one for energy and the other for enstrophy. Since the enstrophy spectrum has larger  $k$  dependency, it tends to get dissipated by viscosity preferentially and as a result the energy spectrum cascades to small wavenumbers and condensates to a minimum wave number that the system allows. This is a process of self-organization of turbulence. The self-organized state can simply be described by variation of minimizing the enstrophy with the constraint of constant energy,

$$\delta(U - \lambda W) = 0 \quad (12)$$

With the help of the definition of the energy and the enstrophy in Eqs (7) and (8), Eq. (12) gives the eigenvalue equation for  $\phi$ ,

$$\nabla^2 \phi + \lambda \phi = 0 \quad (13)$$

Thus may be solved for a given boundary condition and the smallest eigen value gives the self-organized state.

*2.3.4. Weak turbulence theory* If the normalized amplitude of the wave is much smaller than the ratio of the drift wave frequency to the ion cyclotron frequency, the nonlinear term in the Hasegawa-Mima equation may be treated as perturbation and the evolution of the turbulent spectrum may be studied using the weak turbulence theory. We take three waves with wave numbers  $\mathbf{k}_1$ ,  $\mathbf{k}_2$ , and  $\mathbf{k}_3$ , such that

$$k_1^2 \leq k_2^2 \leq k_3^2 \quad (14)$$

The linearized Hasegawa-Mima equation gives the drift wave frequency given by

$$\omega_{\mathbf{k}} = \frac{\mathbf{k} \times \hat{\mathbf{z}} \cdot \nabla \ln(\mathbf{n}_0/\omega_{ci})}{\epsilon(1 + k^2)} \quad (15)$$

and the nonlinear term can be treated as a perturbation to produce coupling of the three waves (Hasegawa et al 1979[11]). In the three wave couplings, the number of wave quanta defined by


$$N_p = (1 + k_p^2)|\phi_p|^2/|k_q^2 - k_r^2|, \quad k_p^2 \neq k_r^2 \quad (16)$$

is conserved so that

$$N_3 - N_1 = const., \quad N_1 + N_2 = const., \quad N_2 + N_3 = const. \quad (17)$$

This result shows that a loss of one quantum of the wave with wave number  $k_2$  appears as a gain in one quantum of the wave with the wave numbers given by  $k_1$  or  $k_3$ . This indicates a dual cascade of energy to smaller and larger wave numbers. A clearer picture that is consistent with the well-known weak turbulence result can be obtained by requiring resonant three wave interactions such that the wave matching condition is satisfied. This is possible in small wave number region such that,

$$k_q^2 - k_r^2 = k_{qy}/\omega_q - k_{ry}/\omega_r = \omega_p M \quad (18)$$



**Figure 1.** Zonal flow pattern predicted from the weak turbulence theory.

where  $k_y$  is the wave number in the direction of  $\hat{z} \times \nabla \ln n_0$  and

$$M \equiv \frac{\omega_p(k_{ry} - k_{qy}) + \omega_q(k_{py} - k_{ry}) + \omega_r(k_{qy} - k_{py})}{3\omega_p\omega_q\omega_r} \quad (19)$$

In this regime, the wave energy decays into smaller frequencies since the wave quanta defined by

$$N_k = W_k / \hbar\omega_k \quad (20)$$

is conserved. Since the drift wave frequency is proportional to  $k_y$  this result indicates that the wave energy decays to that with smaller  $k_y$  keeping  $k_x$  more or less constant at  $1/\rho_s (\equiv k_c)$ . In a cylindrical plasma, this indicates that the wave energy cascades to a mode having zero azimuthal wave numbers as shown in Fig.1, with radial wave number being kept at  $k_c$ . The theory presented by Hasegawa et al (1979)[11] is the first prediction of the formation of the zonal flow.

*2.4. Introduction of the Hasegawa-Wakatani Model and Prediction of the Zonal Flow in Hydrodynamic Regime*

If the free energy source is large enough, the level of wave amplitude will approach to the ratio of the drift wave frequency to the ion cyclotron frequency. Then the drift wave becomes fully nonlinear as indicated by the Hasegawa-Mima equation. This may be the case when the wave is generated by a combination of magnetic field curvature and pressure gradient. Hasegawa and Wakatani (1987) [12] developed a model that describes evolution of the electrostatic potential  $\phi$  and the density fluctuation  $n$  for such a case. They included electron resistivity and ion viscosity, which are responsible to the excitation of the curvature driven instability and dissipation of the excited turbulence. In a cylindrical geometry the equation of vorticity reads,

$$\frac{\rho_s^2}{a^2} \frac{d}{dt} \nabla_{\perp}^2 \phi = (\nabla \ln n \times \nabla \Lambda) \cdot \hat{z} + \frac{\omega_{ce}}{\nu_{ei}} \left( \frac{a}{R} \right)^2 \nabla_{\parallel}^2 (\ln n - \phi) + \frac{\mu}{\omega_{ci} a^2} \nabla_{\perp}^4 \phi \quad (21)$$

Here  $\nabla \Lambda$  represents the curvature of the toroidal magnetic field,  $\omega_{ce}$  and  $\nu_{ei}$  are electron cyclotron frequency and electron-ion collision frequency,  $a$  and  $R$  are minor and major radius of the toroidal container, and  $\mu$  represents the ion viscosity. The equation of continuity is also modified due to the curvature and reads,

$$\frac{d}{dt} \ln n = (\nabla \ln n \times \nabla \Lambda) \cdot \hat{z} + \frac{\omega_{ce}}{\nu_{ei}} \left( \frac{a}{R} \right)^2 \nabla_{\parallel}^2 (\ln n - \phi) \quad (22)$$

Equations (21) and (22) form closed set of nonlinear equations that describe evolution of the potential and density fields. Even in the presence of the curvature terms, it can be shown that in the inviscid limit, the total energy and potential enstrophy are conserved;


$$W = \frac{1}{2} \int dV \left[ (\ln n)^2 + \frac{\rho_s^2}{a^2} (\nabla_{\perp} \phi)^2 \right] \quad (23)$$

and

$$U = \frac{1}{2} \int dV \left[ \frac{\rho_s^2}{a^2} \nabla_{\perp}^2 \phi - \ln n \right]^2 \quad (24)$$

The conservation of  $W$  and  $U$  indicates that the existence of inverse cascade of energy spectrum as discussed earlier. In the case of toroidal plasma, in addition to these two conservations, the total angular momentum in the poloidal direction (which should be zero) should also be conserved. Thus the self-organized state should be obtained by minimizing the potential enstrophy, (23) with constrains of conservation of energy (24) as well as the total toroidal angular momentum. This will lead to a self-organized structure in which the minimum eigen value having a node between the center and the wall. As a result, at self-organization, the radial electric field reverses its sign at some radial position. The minimization of the potential enstrophy with these constraints will lead to generation of global azimuthal flow of the plasma that will have a shear. Equations (21) and (22) are numerically solved to trace the evolution of the turbulence spectrum excited by the magnetic field curvature. Results of the simulation are shown in Fig. 2. Here the left figure is initial potential profile (stream function), which is






**Figure 2.** Equipotential contour of simulation results obtained from Eq. (21) and (22). The solid (dotted) lines are for positive (negative) equipotentials. It is clear that the curvature driven plasma turbulence produces a self-organized state in which a global equipotential that has a  $\phi = 0$  closed surface near outer edge of the plasma.

randomly given, and the right figure shows the potential profile at a final time. Here dotted lines show negative equipotential lines. One can see in this figure that the self-organized state produces equipotential line that has a closed node at some outer radial position indicating formation of a global azimuthal shear flow.

In Fig.3 we show the evolution of the radial potential profile in the simulation. Here the solid line indicates the theoretically predicted self-organized state based on the minimization of the potential enstrophy with constraints of both the total energy and the global angular momentum. This figure shows clearly that the self-organized state produces a global radial electric field that reverses its sign. This electric field creates an  $E \times B$  rotation of plasma in the azimuthal direction that changes direction at some radius indicating a generation of an azimuthal shear flow. This we believe is the first observation of a zonal flow generation in turbulent plasma. The zonal flow structure is global rather than microscopic as indicated in the kinetic theory, Fig.1. It may be due to the fact that the instability here is driven by global magnetic field curvature, which is hydrodynamic.

### *2.5. What are the effects of Zonal Flow on Plasma Transport?*


Zonal flows have been observed to inhibit transport of vortex eddies across the flows. One well-known example is the atmospheric dynamics of the planet Jupiter. As has been demonstrated by Hasegawa et al (1979) [11], the Hasegawa Mima equation has a structure mathematically identical to that which describes the horizontal motion of planet atmosphere with gravity and gradient of Coriolis force. On a planet surface, the latitudinal direction in which the Coriolis parameter varies corresponds to the radial



**Figure 3.** Temporal variations of the radial profile of axis symmetric potential profile are shown for different time. The solid line is the theoretically predicted potential profile obtained from the self-organization theory.

direction in cylindrical plasma, the longitudinal direction corresponds to the azimuthal direction and the vertical direction to the axial direction. Fig.4 is the well-known picture of Jovian atmosphere.


Analogy of the mathematical structure of the model equation that describes the plasma turbulence in the magnetic field to that of the planetary atmospheric motion indicates that a similar turbulent dynamics may operate also in plasmas. The simulation result obtained by Hasegawa and Wakatani (1987) [12] in fact indicate that convection of the plasma vortices seems to be inhibited across the equipotential surface of  $\phi = 0$ .



**Figure 4.** A profile of atmospheric motion on Jovian surface. Existence of longitudinal zonal flows is clearly visible. The zonal flows are seen to inhibit convection of atmospheric vortices in the latitudinal direction crossing the flow.

Since electrons are expected to move on the constant  $\phi$  surface, if vortices cannot convect across this surface, zonal flows are expected to inhibit electron transport in the radial direction. This fact is indicative of a very interesting process in which instability excited self-organized state of plasma may inhibit anomalous electron transport in the radial direction. This contradicts to previously known effect of plasma turbulence that predicts anomalous diffusion. In fact, the reduction of the transport in the presence of zonal flows is verified numerically in resistive pressure-gradient-driven turbulence[13], ion temperature gradient turbulence[14], collisionless trapped electron mode turbulence[15], as well as other models. Fig. 5 shows the reduction of electron heat conduction in the presence of zonal flows.

Although at present existence of zonal flows and their possible role in reduction of anomalous transports have been confirmed in various experiments, theoretical studies



**Figure 5.** Simulation result obtained by Xiao et al (2010) [15]. The figure shows time evolution of electron heat transport. The solid line is the heat transport obtained in the simulation where self-generated zonal flow is present, while the dotted lines is the heat transport in which the zonal flows are artificially removed.

on the stability of the zonal flow and its influence on plasma transport are still actively persuaded. This is the subject of study presented in the next section.

### 3. Zonal Flows and Shear Suppression in Drift Wave Turbulence

In this section, we briefly summarize progress in the theory of zonal flows and velocity shear suppression of drift wave turbulence. The discussion consists of a review of basic zonal flow physics, the dynamics of shearing and its relation to zonal flow formation and the behavior of the dynamical system defined by the coupled flows and fluctuations. Some applications of the theory to problems of current interest are discussed.

A zonal flow[9] is an  $n = 0$ ,  $m = 0$  potential fluctuation with finite  $k_r$ . Given their structure, zonal flows may be thought of as secondary (nonlinearly driven) modes of minimal inertia[10, 11] and minimal damping[16] which drive no transport. Zonal flows are thus natural repositories in which to safely retain energy released by gradient driven

microturbulence.

While there are several approaches to explain zonal flow generations, including modulational calculations via the reductive perturbation method[17, 18, 19, 20], the fundamental mechanism of zonal flow formation in a quasi-2D fluid or plasma is potential vorticity (PV) transport or mixing in a system with one direction of symmetry[21]. Kelvin's theorem underpins this idea. The physics of zonal flow formation was studied within the geophysical fluid dynamics community long before it became a hot topic in the plasma community. Indeed, the quasi-geostrophic equation was first derived by Charney in 1948[22], and within a few years, Charney and Drazin[23] obtained a momentum conservation theorem which elucidated the nonlinear origin and dynamics of zonal flows. In a plasma, vorticity transport or mixing (i.e.  $-\rho^2\langle\tilde{v}_r\nabla^2\tilde{\phi}\rangle$  where  $\langle\dots\rangle$  is an ensemble average) results from the breaking of guiding center ambipolarity and the structure of the gyrokinetic Poisson equation. The flux of vorticity or polarization charge is equivalent to a Reynolds force, as first shown by G.I. Taylor[24], i.e.  $-\rho^2\langle\tilde{v}_{r,E}\nabla^2\tilde{\phi}\rangle = -\partial_r\langle\tilde{v}_{r,E}\tilde{v}_{\perp,E}\rangle$ , where  $\langle\tilde{v}_{r,E}\tilde{v}_{\theta,E}\rangle$  is the Reynolds stress. The first discussion of the relation between vorticity and wave momentum transport and zonal flow formation were by Diamond and Kim[25] and Diamond, et. al.[26]. The fluctuation-driven Reynolds force in turn drives the flow. Vorticity transport depends critically upon the phase relation between  $\tilde{v}_r$  and  $\nabla^2\tilde{\phi}$ . This cross phase is determined by the micro dynamics of the mixing process, i.e. in particular, the velocity shear. There are several viable candidate mixing processes, which include:

- (i) direct dissipation, as by viscosity
- (ii) nonlinear coupling to small scale dissipation by the *forward* cascade of potential vorticity. This effectively replaces the molecular viscosity by an eddy viscosity.
- (iii) Rossby or drift wave absorption at critical layers, where  $\omega = k_x\langle V_x(y)\rangle$ . This is essentially Landau resonance. Transport or mixing of PV region requires the overlap of neighboring critical layers, leading to stochastization of flow streamlines. Stochastization of streamlines guarantees PV mixing much like stochastization of orbits leads to mixing of  $\langle f \rangle$  in quasilinear theory.
- (iv) stochastic nonlinear wave-fluid element scattering, which is analogous to transport induced by nonlinear Landau damping.

Note that the more general concepts are stochasticity of streamlines and forward potential enstrophy cascade to small scale dissipation. Interestingly, when looking at the phenomenon of zonal flow self-organization from the standpoint of PV transport and mixing, it is the *forward enstrophy cascade* which is critical, and not the inverse energy cascade, as is conventionally mentioned!! Finally, we observe that zonal flow acceleration is not necessarily a strongly nonlinear process. PV mixing can occur via wave absorption, and can be manifested in weak turbulence, as an essentially quasilinear process. In this regard, the reader should consult[27]. To relate the transport of PV to the energetics of an open system, it is useful to consider the simplest non-trivial system, namely that of Hasegawa and Wakatani[8]. This system has a conserved inviscid PV

$q = n - \nabla^2 \phi$ . Note the obvious analogy between total PV and total charge (i.e. guiding center + polarization). Conservation of potential enstrophy  $\langle q^2 \rangle$  and some straight forward algebra yield the identity:

$$\langle \tilde{v}_r \nabla^2 \tilde{\phi} \rangle = \langle \tilde{v}_r \tilde{n} \rangle + (\partial_t \langle \tilde{q}^2 \rangle + \partial_r \langle \tilde{v}_r \tilde{q}^2 \rangle + D_0 \langle (\nabla \tilde{q})^2 \rangle) / \langle q \rangle' \quad (25)$$

Here  $\langle \tilde{q}^2 \rangle$  denotes fluctuation potential enstrophy, which was previously defined as  $U$ . This choice is motivated by the consideration that the potential vorticity  $q$  is conserved inviscidly and thus is the fundamental quantity in drift wave - zonal flow dynamics. The prime ( $\langle \dots \rangle'$ ) denotes the radial derivative and the Prandtl number  $Pr \equiv \mu/D_0 = 1$  is assumed for simplicity, where  $D_0$  is the particle diffusivity and  $\mu$  is the viscosity. Equation (25) locks the Reynolds force to the driving flux and the local dissipation of fluctuation potential enstrophy (either by viscosity or local divergence/convergence of the turbulence spreading flux). Using large scale momentum balance, we can extend Equation (25) to derive a zonal flow momentum theorem in the style of a Charney-Drazin theorem [23, 28], i.e.

$$\partial_t \{(\text{GWMD}) + \langle v_\theta \rangle\} = -\langle \tilde{v}_r \tilde{n} \rangle - \delta_t \langle \tilde{q}^2 \rangle / \langle q \rangle' - \nu \langle v_\theta \rangle \quad (26a)$$

$$\text{GWMD} = -\langle \tilde{q}^2 \rangle / \langle q \rangle' \quad (26b)$$

Here GWMD is the generalized wave momentum density (i.e. pseudomomentum) and  $\delta_t \langle \tilde{q}^2 \rangle / \langle q \rangle' \equiv (\partial_r \langle \tilde{v}_r \tilde{q}^2 \rangle + D_0 \langle (\nabla \tilde{q})^2 \rangle) / \langle q \rangle'$ . Equation (26a) states that in the absence of a driving flux or a local potential enstrophy decrement, it is impossible to accelerate or maintain a zonal flow against drag with time stationary fluctuations. Physically, Equation (26a) expresses the freezing in of quasi-particles into the flow, unless 'slippage' due to drive or dissipation is induced. Equation (26a) imposes a fundamental constraint on models of stationary zonal flows. These *must* be constructed using an explicit connection to the driving turbulent flux and fluctuation dissipation. Note that flow acceleration requires relative spatial separation of fluctuation drive and damping, as in drift wave eigenfunctions with outgoing wave structure [25].

Of course, zonal flows are of great interest in fusion due to the fact that they can regulate turbulence by shearing. Fig.6 illustrates the effect of a self-generated shear flow on extended turbulent eddys. This effect was first explicitly stated and developed in the seminal paper by Biglari, Diamond and Terry in 1990 [30]. Time stationary, coherent shearing has long been known to couple to diffusive scattering to yield an enhanced, hybrid decorrelation rate. This process was discussed by Kelvin, G.I. Taylor [24], Dupree [29], and then applied to tokamak shear flows by Biglari, Diamond and Terry. The hybrid decorrelation rate is

$$\frac{1}{\tau_{c,\mathbf{k}}} = \left( \frac{k_\theta^2 \langle v_E \rangle'^2 D_\perp}{3} \right)^{1/3} \quad (27)$$

When  $k_\theta \langle v_E \rangle' \Delta_r > D_\perp / \Delta_r^2$ , decorrelation will be enhanced relative to the rate for zero shear. Important corrections to the theory of shear decorrelation related to shaping and compression were noted by Hahn and Burrell [31]. Other, linear shearing effects include

spatial resonance dispersion ( $\omega - k_{\parallel}v_{\parallel} \rightarrow \omega - k_{\parallel}v_{\parallel} - k_{\theta}\langle v_E \rangle'(r - r_0)$ ) and differential response rotation. An important caveat is that modes can adjust to weaken the effects of externally prescribed shear[32]. Thus, there is a qualitative difference between the effects of self-consistent, energy conserving zonal shears and externally prescribed shears. Zonal shears are most effectively described by wave kinetics, based on eikonal theory[9, 33]. In wave kinetics, shear acts to tilt and stretch vortices, thus increasing  $k_r$ . For a coherent shear, we have  $k_r = k_r^{(0)} - k_{\theta}\langle v_E \rangle'\tau$  after a time duration  $\tau$ , while for wave packet propagation in a complex shearing pattern, the wave packet undergoes induced diffusion in  $k_r$ , so  $\langle \delta k_r^2 \rangle = D_k \tau$ , where  $D_k = \sum_q k_{\theta}^2 |\tilde{v}_E, q'|^2 \tau_{k,q}$ . The physics of induced diffusion is a random walk of the wave packet  $k_r$  due to wave ray chaos, caused by overlap of resonances between zonal shear phase velocities and wave packet group velocity. Here  $\tau_{k,q}$  is the correlation time of a wave packet  $\mathbf{k}$  with zonal mode  $\mathbf{q}$ . The mean wave action (population) density equation thus becomes:


$$\frac{\partial}{\partial t} \langle N \rangle - \frac{\partial}{\partial k_r} D_{\mathbf{k}} \frac{\partial \langle N \rangle}{\partial k_r} = \gamma_k \langle N \rangle - \langle C\{N\} \rangle \quad (28)$$

As a consequence, the effect of zonal flow shearing on wave energy is given by


$$\frac{\partial}{\partial t} \langle \epsilon \rangle = - \int d\mathbf{k} v_{gr}(\mathbf{k}) D_{\mathbf{k}} \frac{\partial \langle N \rangle}{\partial k_r} \quad (29)$$

so that for wave enstrophy density decreasing with  $k_r$  (i.e.  $d\langle \Omega \rangle / dk_r < 0$ ), wave energy is damped. The fate of the energy is clear - it goes to the zonal flow via Reynolds work, as can be demonstrated by a modulational instability calculation[9, 34]. The bottom line, then, is that for zonal flows “Reynolds work” and “flow shearing” emerge as relabelings of one another, so that the energy budget always balances, intrinsically. This relation is the energy counterpart of the Charney-Drazin theorem for momentum[28] and of the proof of conservation of energy between waves (zonal modes) and particles (drift wave packets) in quasi-linear theory. In this light, the zonal flow damping mechanism emerges as critical to the self-regulation process, as it is a damping for the whole *system*. This follows from the structure of the predator-prey dynamics paradigm applicable to the drift wave-zonal flow interaction.

We have established that zonal flows regulate drift waves by shearing, while at the same time drift wave turbulence pumps the zonal flows. These two populations thus naturally form a self-regulating ‘predator-prey’ feedback loop. This loop is depicted in Fig.8. The control parameters for this system are the sources (i.e. heat flux, etc.) and the zonal flow damping. Interestingly, the predator-prey system reflects the underlying dual cascade of quasi-2D fluids. To see this, note that the prey equation for mean wave action density (related to wave potential enstrophy density) involves induced diffusion to high  $k_r$ , which is analogous to the forward enstrophy cascade. Likewise, the prey equation for the zonal flow energy describes the ‘negative viscosity phenomenon’ of zonal flow growth by wave Reynolds work and so is clearly analogous to the inverse energy cascade. We emphasize, however, that these two transfer processes are *not* local, as in the conventional cascade picture. The mean field predator-prey system may



**Figure 6.** Flow effect on fluctuation. This figure, from ref.[36], illustrates the effect of shearing on fluctuation structure and scale.



**Figure 7.** Stretching of eddy

be reduced to a simple dynamical system,

$$\partial_t N = \gamma N - \alpha V^2 N - \Delta \omega N^2 \quad (30a)$$

$$\partial_t V^2 = \alpha N V^2 - \gamma_d V^2 - \alpha_2 V^4 \quad (30b)$$


Here  $\gamma$  is the drift wave growth rate,  $\Delta \omega$  is the turbulence nonlinear decorrelation rate,  $\gamma_d$  is the frictional zonal flow damping and  $\alpha$ ,  $\alpha_2$  are coefficients in the model, as given by Eqs.30a and 30b and discussed in ref.[9]. This system exhibits multiple



fixed points corresponding to no-flow and finite-flow states, respectively. These are discussed in [9, 35]. The roots and states are tabulated in Fig.9. A value of the local growth, operationally set by the heat flux, sets the transition threshold, and the state transition occurs by a super-critical bifurcation in which a relaxation mode softens and goes unstable. Simple numerical experiments[36] have verified the predictions of this model(Fig.10). In particular, these demonstrate the important role of zonal flow damping in regulating the drift wave-zonal flow system - see Fig.11. This system, while conceptually instructive, is over-idealized. To address realistic problems, such as the  $L \rightarrow H$  and ITB transition, models with multiple predators, corresponding to zonal and mean shear flows, have been developed. Here, ‘mean flows’ refer to mean field  $\langle \mathbf{E} \rangle \times \mathbf{B}$  shear flows, where the radial electric field  $\langle E_r \rangle$  must satisfy the radial force balance equation. Thus, mean flows vary in space on profile scales. In contrast, zonal flows can occur on all meso scales  $l$ ,  $\Delta_c < l < L_p$  where  $L_p$  is a profile scale length and  $\Delta_c$  is the turbulence correlation length. Zonal flow spatial scales are set by those of the turbulence fluctuation intensity. The two predator-one prey model of Kim and Diamond[37], introduces such competition between mean and zonal shears, as well as a link between growth and mean shear on account of the *common* role of the pressure gradient in determining each. In this scenario, the zonal flow triggers the transition while the mean diamagnetic flow locks it in. Note that the concept of local gradient transport bifurcations builds upon the work of Itoh and Itoh[38], and F.L. Hinton[39]. The system evolution as predicted by this models is shown in Fig.12. A slow power ramp reveals a period of dithering or limit cycle oscillations as the transition threshold is approached. During this period, turbulent fluctuations, zonal flows, and  $\nabla p$  all oscillate with a heat flux dependent relative phase. The competition between mean and zonal shears leaves a distinct signature upon the evolving wave form. In the fully evolved H-mode, turbulence is heavily suppressed. Thus, the zonal flow collapses, leaving only the mean flow to support the transport barrier. The multi-predator models (Malkov and Diamond[40]) exhibit hysteresis (even in 0D!). Hysteresis is stronger in 1D models. Note that in these multi predator-prey models, the zonal flow controls access to the intermediate phase while the mean flow ‘locks in’ the H-phase.

Here, an interesting question arises, concerning the interplay of mean electric fields (n.b. we include neoclassical fields in the label ‘mean’) and zonal flows. The evolution of the sheared neoclassical electric field has been studied extensively[41] and recently has been included in full- $f$ , global gyrokinetic simulations[42]. It is important to note that, as stated in ref.[37], the mean and zonal electric shears are *not* simply additive, but instead have a rather complex network of feedback loops. In the multi-predator prey model:

- (i) the pressure gradient can contribute to both prey and predator populations - the former via, say, ITG mode growth (i.e.  $\gamma = \gamma(\nabla p)$ ) and the latter by its contribution to mean shears via radial force balance.
- (ii) mean field shear has a strong effect on the cross-phase of the fluctuation Reynolds



**Figure 8.** Predator-prey feedback loop structure[9] for the drift wave-zonal flow system, i.e. Eqs. (30a) and (30b).

stress  $\langle \tilde{v}_r \tilde{v}_\theta \rangle$ , which drives the zonal flow. As a consequence, mean field shear enters zonal flow dynamics *multiplicatively* and *nonlinearly*, so mean field shear can effectively reduce and/or cut off the generation of zonal flows by potential vorticity transport or inverse cascade. This is part of the  $L - H$  transition model proposed in ref.[37].

- (iii) zonal flows can, of course, enhance the growth of the mean shear by extracting fluctuation energy, thus reducing transport and so allowing  $\nabla p$  to steepen, as discussed in ref.[37].
- (iv) the mean electric field directly enters the zonal flow dielectric [16, 43].

Indeed, the interaction of mean (neoclassical) and zonal shears is especially interesting in the context of stellarators, where purely neoclassical electric field bifurcations are possible and additional control parameters are available. All told, while the interaction of zonal and mean flows has been studied extensively, much work is still required for a complete understanding.


The basic ideas of zonal flow physics discussed here have been applied to numerous

State	No flow	Flow ( $\alpha_2 = 0$ )	Flow ( $\alpha_2 \neq 0$ )
$N$ (drift wave turbulence level)	$\frac{\gamma}{\Delta\omega}$	$\frac{\gamma_d}{\alpha}$	$\frac{\gamma_d + \alpha_2\gamma\alpha^{-1}}{\alpha + \Delta\omega\alpha_2\alpha^{-1}}$
$V^2$ (mean square flow)	0	$\frac{\gamma}{\alpha} - \frac{\Delta\omega\gamma_d}{\alpha^2}$	$\frac{\gamma - \Delta\omega\gamma_d\alpha^{-1}}{\alpha + \Delta\omega\alpha_2\alpha^{-1}}$
Drive/excitation mechanism	Linear growth	Linear growth	Linear growth Nonlinear damping of flow
Regulation/inhibition mechanism	Self-interaction of turbulence	Random shearing, self-interaction	Random shearing, self-interaction
Branching ratio $\frac{V^2}{N}$	0	$\frac{\gamma - \Delta\omega\gamma_d\alpha^{-1}}{\gamma_d}$	$\frac{\gamma - \Delta\omega\gamma_d\alpha^{-1}}{\gamma_d + \alpha_2\gamma\alpha^{-1}}$
Threshold (without noise)	$\gamma > 0$	$\gamma > \Delta\omega\gamma_d\alpha^{-1}$	$\gamma > \Delta\omega\gamma_d\alpha^{-1}$

**Figure 9.** Roots of predator-prey system. This, from [9], gives the various roots of the multi-state predators-prey system. Note that both no flow and finite flow states are possible.

problems of current interest, which include but are not limited to: poloidal rotation[44], turbulence spreading[45, 46], confinement in plasmas with RMP[47, 48] and beta-plane MHD models of the solar tachocline[49, 50]. The interested reader is referred to the literature, as space does not permit even a short discussion of these. Rather we briefly discuss the current status of our understanding of flow shear dynamics in the  $L \rightarrow H$  transition.

The  $L \rightarrow H$  transition problem is singled out because since its discovery in 1982[51], the  $L \rightarrow H$  transition has driven much of the thinking and research on flows in magnetic fusion. In some sense, it has become *the* classic paradigm problem on flows in tokamaks. Also, the past 2 years have witnessed a remarkable spurt of progress in diagnosing flow and fluctuation behavior in edge plasmas at the transition threshold. In particular, studies by Conway, et.al. on AUG[52], Estrada and Hidalgo[53, 54] on TJ-II, Schmitz[55] on DIII-D, McKee and Z. Yan on DIII-D[56, 57], Kamiya and Ida[58] on JT-60U, G.S. Xu[59] on East, A.Hubbard[60] on Alcator C-Mod, and S.J. Zweben[61] on NSTX stand out as significant contributions. The gist of this wealth of information is that for  $P \sim P_{th}$ , limit cycle or dithering oscillations are observed in the flows and turbulence. These cyclic phenomena characterize what is now called the I-phase (I for intermediate, between L and H),. Multi-Shear flow competition is at work in the I-phase, and the flow structure evolves as the transition progresses. Many aspects of the observed phenomena are consistent with the multi-predator shearing model[37]. In particular, the variation of relative phases with increasing heat flux is observed. These




**Figure 10.** Time evolution of fluctuation intensity and turbulence driven flow from the simulation of ref.[36]

are a variety of suggestions or hints as to the trigger mechanism. These include the GAM, zonal flow, mean  $\mathbf{E} \times \mathbf{B}_0$  flow, mean poloidal flow and more. It is not at all obvious to this author that there need be a unique route to the transitions. Different shear flows may act as trigger under different circumstances. To face this challenge, theory should forsake the 0D models for one space-one time dimensional models of edge layer evolution. Theory should *predict* something qualitatively new. One suggestions as to where to look is the flow shear and fluctuation evolution during a slow, ELM free back-transition. Also, theory should link micro-dynamics to macroscopics, such as the power threshold scalings. Both theory and experiment should elucidate SOL flow effects[62] on the shear structure *inside* the separatrix. This important issue remains terra incognita. Finally, basic experiments should be confronted to elucidate the details of the microscopic physical processes involved in shear layer formation[63]. Readers should stay tuned for excitement in the near future.

#### 4. Inverse cascade and whistlerization of self-generated low frequency electro-magnetic fluctuation: An extension of Hasegawa-Mima equation

In this section, the nonlinear dynamics of electron magneto hydrodynamics (EMHD) is reviewed. The EMHD plays an important role in space and magnetic confinement




**Figure 11.** Fluctuation and flow damping. This, from a simulation of negative compressibility turbulence presented in ref.[36], presents the scaling of the saturated fluctuation intensity ( $\langle (\tilde{n}/n_0)^2 \rangle$ ) with zonal flow damping  $\mu$ . Note for small or moderate damping, the saturated intensity scales linearly with  $\mu$ , as predicted by the predator-prey model. For strong damping, the zonal flow is effectively suppressed, so  $\langle (\tilde{n}/n_0)^2 \rangle$  becomes independent of  $\mu$ .

plasmas, in particular, in collisionless reconnection and in relativistic laser plasmas produced by ultra intense short pulse laser, in particular, in the relativistic electron transport. In laser plasmas, magnetic fields are generated by the Weibel instability related to the relativistic electron transport which has been widely investigated in the fast ignition research. In this section, the inverse cascade and the whistlerization in the Weibel turbulence are discussed in some detail.

#### 4.1. Basic equations

The low frequency electro-magnetic fluctuations (whistler turbulence) are generated by relativistic electron beams ( electro-magnetic two stream instability ) and/or by non-isotropy of electron momentum distribution (Weibel instability). When the scale length or the time scale of fluctuation is much shorter or faster than the ion scales, ions are assumed immobile and the incompressible electron fluid dynamics is taken into account. Namely, nonlinear dynamics of the fluctuation is described by the electron hydro dynamics (EMHD). The basic equations of EMHD are; The magnetic field



**Figure 12.** 2 predators and 1 prey. This figure, from ref.[37], shows the evolution of the turbulence intensity, zonal flow intensity, and mean pressure gradient during a power ramp through an  $L \rightarrow H$  transition. Here the solid line is the fluctuation (drift wave) intensity, the dotted line is the zonal flow intensity, and the dashed line is the mean pressure gradient. The  $x$  axis is normalized heat flux  $Q$ . Note that as heat flux  $Q$  increases, the system first enters a cyclic state, where fluctuation intensity and zonal flow intensity oscillate, while  $\nabla p$  grows very slowly. At sufficient  $Q$ , turbulence and zonal flow collapse, and  $\nabla p$  increases rapidly. This final state corresponds to an H-mode.

induction equation:

$$\partial_t \mathbf{B} = -\nabla \times \mathbf{E} \quad (31)$$

Electron fluid equation of motion:

$$\partial_t \mathbf{p} + \mathbf{v} \cdot \nabla \mathbf{p} = -e(\mathbf{E} + \mathbf{v} \times \mathbf{B}) - \frac{1}{n} \nabla P - \nu \mathbf{p} \quad (32)$$

Ampere's law:

$$\frac{\nabla \times \mathbf{B}}{\mu} = \mathbf{j} = -en_0 \mathbf{v} + \mathbf{j}_{ex} \quad (33)$$

Here, ions are immobile back ground and the electron fluid is assumed incompressible and the external current,  $\mathbf{j}_{ex}$  represents current carried by beam and/or high energy electron.

By introducing generalized vorticity,  $\mathbf{\Omega} = \mathbf{\Omega}_B - \nabla \times \mathbf{v}$ , Eqs. (31) and (32) yield the vorticity equation;

$$\partial_t \mathbf{\Omega} - \nabla \times (\mathbf{v} \times \mathbf{\Omega}) = \nabla \times \left( \frac{\nabla P}{mn} + \nu_e \mathbf{v} \right) \quad (34)$$

The right hand side of equation (34) is the source (thermo electric effect ) and dissipation of the vorticity. When the thermoelectric effects and the external current do not exist, the equation (33) and (34) are combined to obtain a magnetic field evolution equation;

$$\partial_t \mathbf{\Omega} - \nabla \times (\mathbf{v} \times \mathbf{\Omega}) = \nabla \times \nu_e \mathbf{v} \quad (35)$$

$$\mathbf{\Omega} = \mathbf{\Omega}_B - \frac{c^2}{\omega_p^2} \nabla^2 \mathbf{\Omega}_B \quad (36)$$

$$\mathbf{v} = -\frac{c^2}{\omega_p^2} \nabla \times \mathbf{\Omega}_B \quad (37)$$

where  $\mathbf{\Omega}_B = e\mathbf{B}/m$  is the electron cyclotron frequency. In the equation (36), the second term is finite only when the magnetic fluctuation scale length is the order of electron skin depth. This is the electron Hall effect and the set of equations (35),(36), and (37) are the Hall EMHD equations. For the Weibel instability, beam instability or magnetic reconnection, the turbulence is essentially two dimensional like the drift wave turbulence. In the two dimensional turbulence, it is possible to introduce two scalar quantities,  $\psi$  and  $b$  and represent  $\mathbf{\Omega}_B = e\mathbf{B}/m$  as

$$\mathbf{\Omega}_B = \hat{z} \times \nabla \psi + \hat{z} b \quad (38)$$

Note that  $\psi$  is the magnetic flux function and the equi-contours of  $\psi$  represent magnetic field lines. The equations for  $\psi$  and  $b$  are obtained from the Eqs. (35),(36), and (37) by normalizing the space variables by the electron skin depth as follows,


$$\partial_t (\psi - \nabla^2 \psi) + \hat{z} \times \nabla b \cdot \nabla (\psi - \nabla^2 \psi) = \eta \nabla^2 \psi \quad (39)$$

$$\partial_t (b - \nabla^2 b) - \hat{z} \times \nabla b \cdot \nabla \nabla^2 b + \hat{z} \times \nabla \psi \cdot \nabla \nabla^2 \psi = \eta \nabla^2 b \quad (40)$$

Here, the right hand side of the Eqs. (39) and (40) is the magnetic field dissipation due to resistivity. These equations are derived in [75]. It is known that the turbulence described by the Eqs.(39) and (40) exhibits inverse cascade or beam current filament merging. Note also that the first two terms of the Eq. (40) are the same as Hasegawa-Mima equation and the other nonlinear terms have similar structure, namely, vector nonlinearity. From those characters of the equations, it is predicted that the inverse cascade occurs.

#### *4.2. An example of nonlinear evolution of EMHD turbulence and self-organization in laser plasmas*

The quasi-static magnetic field fluctuations are generated in relativistic laser plasmas. The Fig.13(a) shows that an intense laser irradiates the over-dense from the left hand side and relativistic electrons are generated and propagate from the left to right. Magnetic fields perpendicular to the simulation plane are generated and amplified. In



**Figure 13.** Quasi-static magnetic field generation by transverse two stream / Weibel instability by intense electron beam. (a) 2-D PIC simulation for the laser interaction with over dense plasma (5 times cut-off density). Magnetic field generation by the instability in laser produced plasmas, (b) 2-D Hybrid simulation: particle (electron beam) and fluid (background plasma with the density of 10 times electron beam density). The electron beam density is shown in the figure. This figure indicates beam driven transverse two stream instability and merging of electron beam.


the Fig.13(b), initially a relativistic beam is injected perpendicular to the simulation plane and temporal evolution of beam density distribution in the transverse plane is shown. This indicates that the electron beam break up into filaments and then they merged into few filaments. In the following, the merging process of electron beam filaments is discussed as the inverse cascade in the low frequency magnetic field turbulence. The Fig.14 [68] shows self generated magnetic field temporal evolution, where the center of the initial beam is at the upper right corner. At the beginning, cylindrically symmetric magnetic field is generated due to the background electron shear flow.

Assuming the total vorticity of back ground electron is conserved, then,

$$(\hat{z} \times \nabla(\psi - \nabla^2\psi))_\theta \sim \partial_r(\psi - \partial_r^2\psi) = A\delta(r - r_0) \quad (41)$$

The solution of the Eq. (41) is easily obtained to show that the  $B$  field is localized and reversed at  $r = r_0$  with the thickness of the order of the skin depth. This initial cylindrically symmetric magnetic field structure is consistent with the Fig.13-a. The cylindrical symmetry of the magnetic field is broken by the tearing instability.





**Figure 14.** 2-D Hybrid simulation where a relativistic electron beam propagates perpendicular to the X-Y plane. Temporal evolution of magnetic field lines excited by an electron beam (electron beam density is 10% of the back ground electron density) at  $\omega_p t = 67$  and (b)  $\omega_p t = 81$ . The arrows in the figures show total electron fluxes.

Let us take a initial magnetic field as

$$\psi_0 = A \exp(ik_0 x) + c.c. \quad (42)$$

and take perturbed magnetic fields as

$$\psi_1 = g \exp(iky) \text{ and } b = h \exp(ik_0 x +iky) \quad (43)$$

This mode structure is essentially the same as that of the Fig.13-b. The growth rate can be obtained by the linearized equation for  $\psi_1$  and  $b$ ;

$$\partial_t(\psi_1 - \nabla^2 \psi_1) + \hat{z} \times \nabla b \cdot \nabla(\psi_0 - \nabla^2 \psi_0) = 0 \quad (44)$$

$$\partial_t(b - \nabla^2 b) + \hat{z} \times \nabla \psi_1 \cdot \nabla \nabla^2 \psi_0 = 0 \quad (45)$$

The growth rate of this mode is given as follows.


$$\gamma(k) = B_0 k k_0 \left\{ \frac{(1 + k_0^2)(k_0^2 - k^2)}{(1 + k_0^2 + k^2)(1 + k^2)} \right\}^{1/2} \quad (46)$$

This result indicates that both shorter and longer wavelength magnetic field fluctuations are simultaneously excited. The ratio of the energy of longer wavelength mode to that of the shorter wavelength mode is,

$$\frac{(1 + k^2)B_{\perp}^2}{(1 + k_0^2 + k^2)B_z^2} = \frac{k^2(1 + k_0^2)}{k_0^2 - k^2} \quad (47)$$

This ratio is greater than unity for the maximum growing mode. This indicates the inverse cascade of the two dimensional EMHD is similar to the case of Hasegawa-Mima equation [79]. The corresponding growth rate for the Hasegawa-Mima equation is

$$\gamma(k) = A k^2 k_0 \left\{ \frac{k_0^2 - k^2}{(1 + k_0^2 + k^2)(1 + k^2)} \right\}^{1/2} \quad (48)$$



**Figure 15.** Maximum magnetic field strength in the simulation region. Here,  $B_T$  is the transverse component to the beam propagation, and  $B_z$  is the parallel components to it.

In the simulation of [68],  $k_0 > 1.0$ , namely, the wavelength is comparable to the skin depth,  $c/\omega_p$ , then the maximum growth rate is  $\gamma(k) = 1.2\omega_c$  for  $kc/\omega_p = 1.5$ ,  $k_0c/\omega_p = 3$ . According to the simulation  $\gamma \sim 0.24\omega_p$  for  $\omega_c \sim 0.2\omega_p$ .

After the long time inverse cascade,  $k_0^2, k^2 \ll 1$  and  $\gamma(k) \sim \omega_c(kc/\omega_p)^2$  is the time scale of long wave length whistler wave and  $(B_z/B_\perp)^2$  will be 1 for the fastest growing mode according to Eqs. (46) and (48). The Fig.15 [68] shows that the nonlinear evolution of beam driven magnetic fluctuations can be reasonably described by the background electron nonlinearity, although high energy electron beams also respond to the magnetic field fluctuations.

As the final state, it is expected that magnetic bubbles of helical structure are formed in the two dimensional EMHD turbulence. The equations (39) and (40) have three conserved quantities, which are energy, magnetic flux, and cross helicity as derived by A. Das et. al. [75]. Minimizing the energy for a fixed flux and cross helicity, an attractor of magnetic structure evolution can be obtained. It will be a helical structure with  $\langle B_z \rangle^2 = \langle B_\perp \rangle^2$ . Further analysis will be continued in future studies.

## 5. Conclusion

As to conclusions, there are none. This topic is alive and well, and will evolve dynamically. The cross-disciplinary dialogue with the geophysical and astrophysical fluids has surely been beneficial and should continue. Indeed, we still have much to learn. As theorists should *predict*, rather than merely explain, here we predict that this will not be the last prize awarded for the theory of drift wave-zonal flow turbulence.

## Acknowledgments

The authors would like to express their gratitude to EPS for providing the opportunity of submitting this manuscript in connection to the Alfvén Prize award. A. Hasegawa is grateful to late Professor Toshiya Taniuti who guided him to the realm of nonlinear physics, in particular to soliton and vortex dynamics. He is indebted to late Professor Masahiro Wakatani who provided him to continued opportunities to work together in Kyoto and performed critical computer simulations of turbulent plasmas. A. Hasegawa is also grateful to Prof. Zhihong Lin for valuable discussions and providing him critical simulation results related to zonal flow generations.

One of the authors (PHD) acknowledges the superb scientific and human mentorship by the late Marshall N. Rosenbluth. PHD thanks Thomas H. Dupree, B.A. Carreras, Sanae-I. Itoh, Kimitaka Itoh, Taik Soo Hahm and Ozgur Gurcan for their valuable contributions to these ideas and work at various points in his career. PHD also thanks many other theoretical and experimental colleagues and co-workers for their contributions. This work was supported by the United States D.O.E and the Korean N.R.F. W.C.I. Program. Finally, PHD adds that from 2001 to the present, the biannual Festival de Theorie, organized by the C.E.A. of France, has been a valuable source of scientific stimuli.

All the authors thank Yusuke Kosuga for valuable assistance with the preparation of this manuscript.

## References

- [1] R.E. Slusher and C.M. Surko, *Phys. Rev. Lett.* **37**, 1747, (1976).
- [2] E. Mazzucato, *Phys. Rev. Lett.* **37**, 792, (1976).
- [3] R.E. Slusher and C.M. Surko, *Phys. Rev. Lett.* **40**, 400, (1978).
- [4] R.H. Kraichnan, *J. Fluid Mech* **59**, 745, (1973).
- [5] A. Hasegawa and K. Mima, *Phys. Fluids* **21**, 1, (1978).
- [6] L. Onsager, *Supple to Nuovo Cim.* **6**, 279, (1949).
- [7] A.N. Kolmogorov, *Dokl. Akad. Nauk SSR* **30**, 301, (1941).
- [8] A. Hasegawa and M. Wakatani, *Phys. Rev. Lett.* **50**, 682 (1983).
- [9] P.H. Diamond, S.-I. Itoh, K. Itoh, and T.S. Hahm, *Plasma Phys. Control. Fusion* **47**, R35, (2005).
- [10] R.Z. Sagdeev, V.D. Shapiro, and V.I. Shevchenko, *Sov. J. Plasma Phys.* **4**, 306 (1978).
- [11] A. Hasegawa, C.G. MacLennan, and Y. Kodama, *Phys. Fluids* **22**, 2122 (1979).
- [12] A. Hasegawa and M. Wakatani, *Phys. Rev. Lett.* **59**, 1581 (1987).
- [13] B.A. Carreras, V.E. Lynch, and L. Garcia, *Phys. Fluids B* **3**, 1438 (1991).

- [14] R.E. Waltz, G.D. Kerbel, and J. Milovich, *Phys. Plasmas* **1**, 2229 (1994).
- [15] Y. Xiao, I. Holad, W. Zhang, S. Klasky, and Z. Lin, *Phys. Plasmas* **17**, 022302 (2010).
- [16] M.N. Rosenbluth and F.L. Hinton, *Phys. Rev. Lett.* **80**, 724 (1998).
- [17] S. Champeaux and P.H. Diamond, *Phys. Lett. A* **288**, 214 (2001)
- [18] T. Taniuti and N. Yajima, *J. Math. Phys.* **10**, 1369 (1969).
- [19] K. Nozaki, T. Taniuti, and K. Watanabe, *J. Phys. Soc. Japan* **46**, 983 (1979)
- [20] K. Nozaki, T. Taniuti, and K. Watanabe, *J. Phys. Soc. Japan* **46**, 991 (1979)
- [21] D.G. Andrews and M.E. McIntyre, *J. Fluid Mech.* **89**, 647 (1978).
- [22] J.G. Charney, *Geof. Publ.* **17**, 3 (1948).
- [23] J.G. Charney and P.G. Drazin, *J. Geophys. Res.* **66**, 83 (1961).
- [24] G.I. Taylor, *Phil. Trans. Roy. Soc. London.* **A215**, 1 (1915).
- [25] P.H. Diamond and Y.B. Kim, *Phys. Fluids B* **5**, 2343 (1991).
- [26] P.H. Diamond, et.al., *Plasma Physics and Controlled Nuclear Fusion Research 1992* **2**, IAEA, (Vienna), (1993).
- [27] S.M. Tobias, K. Dagon, and J.B. Marston, *ApJ* **727**, 127 (2011).
- [28] P.H. Diamond, O.D. Gurcan, T.S. Hahm, K. Miki, Y. Kosuga, and X. Garbet, *Plasma Phys. Control. Fusion* **50**, 124018 (2008)
- [29] T.H. Dupree, *Phys. Fluids* **9**, 1773 (1966).
- [30] H. Biglari, P.H. Diamond, and P.W. Terry, *Phys. Fluids B* **2**, 1 (1990).
- [31] T.S. Hahm and K.H. Burrell, *Phys. Plasmas* **2**, 1648 (1995).
- [32] B.A. Carreras, K. Sidikman, P.H. Diamond, P.W. Terry, and L. Garcia, *Phys. Fluids B* **4**, 3115 (1990).
- [33] P.H. Diamond, S. Champeaux, M. Malkov, A. Das, I. Gruzinov, M.N. Rosenbluth, C. Holland, B. Wecht, A.I. Smolyakov, F.L. Hinton, Z. Lin, and T.S. Hahm, *Nucl. Fusion* **41**, 1067 (2001).
- [34] L. Chen, Z. Lin, and R. White, *Phys. Plasmas* **7**, 3129 (2000).
- [35] P.H. Diamond, Y.-M. Liang, B.A. Carreras, and P.W. Terry, *Phys. Rev. Lett.* **72**, 2565 (1994).
- [36] L.A. Charlton, B.A. Carreras, V.E. Lynch, K.L. Sidikman, and P.H. Diamond, *Phys. Plasmas* **1**, 2700 (1994).
- [37] E.-J. Kim and P.H. Diamond, *Phys. Rev. Lett.* **90**, 185006 (2003).
- [38] S.-I. Itoh and K. Itoh, *Phys. Rev. Lett.* **60**, 2276 (1988).
- [39] F.L. Hinton, *Phys. Fluids B* **3**, 696 (1991).
- [40] M.A. Malkov and P.H. Diamond, *Phys. Plasmas* **16**, 012504 (2009).
- [41] V. Rozhansky and M. Tendler, *Phys. Fluids B* **4**, 1877 (1992).
- [42] C.S. Chang, S. Ku, P.H. Diamond, Z. Lin, S. Parker, T.S. Hahm, and N. Samatova, *Phys. Plasmas* **16**, 056108 (2009).
- [43] H. Sugama and T.-H. Watanabe, *Phys. Plasmas* **13**, 012501 (2006).
- [44] C.J. McDevitt, P.H. Diamond, O.D. Gurcan, and T.S. Hahm, *Phys. Plasmas* **17**, 112509 (2010).
- [45] O.D. Gurcan, P.H. Diamond, T.S. Hahm, and Z. Lin, *Phys. Plasmas* **12**, 032303 (2005).
- [46] O.D. Gurcan, P.H. Diamond, and T.S. Hahm, *Phys. Plasmas* **13**, 052306 (2006).
- [47] M. Leconte and P.H. Diamond, in press, *Phys. Plasmas*, (2011).
- [48] Y. Xu, et.al., *Nucl. Fusion* **51**, 063020 (2011).
- [49] D.W. Hughes, R. Rosner, and N.O. Weiss, *The Solar Tachocline*, Cambridge University Press (Cambridge), (2007).
- [50] S.M. Tobias, D.W. Hughes, and P.H. Diamond, *ApJ* **667**, L113 (2007).
- [51] F. Wagner, et.al., *Phys. Rev. Lett.* **49**, 1408 (1982).
- [52] G.D. Conway, et.al., *Phys. Rev. Lett.* **106**, 065001 (2011).
- [53] T. Estrada, et.al., *Plasma Phys. Control. Fusion* **51** 124015 (2009)
- [54] T. Estrada, et.al., *EPL* **92**, 35001 (2010).
- [55] L. Schmitz, et.al., submitted to *Phys. Rev. Lett.* , (2011).
- [56] G.R. McKee, et.al., *Nucl. Fusion* **41**, 1235 (2001).
- [57] Z. Yan, et.al., *Phys. Plasmas* **18**, 056117 (2011).

- [58] K. Kamiya, et.al., *Phys. Rev. Lett.* **105**, 045004, (2010).
- [59] G.S. Xu, et.al., submitted to *Phys. Rev. Lett.* , (2011).
- [60] A.E. Hubbard, et.al., *Phys. Plasmas* **18**, 056115 (2011).
- [61] S.J. Zweben, et.al., *Phys. Plasmas* **17**, 102502 (2010).
- [62] B. LaBombard, et.al., *Nucl. Fusion* **44**, 1047 (2004).
- [63] M. Xu, et.al., *Phys. Rev. Lett.* , (2011), in press
- [64] A.S. Kingsep, K.V. Chukbar, and V.V. Yankov, *Reviews of Plasma Physics Vol. 16*, Consultant Bureau (New York), (1990). *Reviews of Plasma Physics*, 1408 (1982).
- [65] M. Honda, J. Myer-ter-Vehn, and A. Pukov, *Phys. Plasmas* **7**, 1302 (2000).
- [66] M. Honda, J. Myer-ter-Vehn, and A. Pukov, *Phys. Rev. Lett.* **85** 2128 (2000)
- [67] Y. Sentoku, K. Mima, P.K. Kaw, and K. Nishikawa, *Phys. Rev. Lett.* **90**, 155001 (2003).
- [68] T. Taguchi, T.M. Anonsen,Jr., C.S. Liu, and K. Mima, submitted to *Phys. Rev. Lett.* **86** 5055 (2001).
- [69] H.B. Cai, S.P. Zhu, M. Chen, S. Wu, X.T. He, and K. Mima, *Phys. Rev. E* **83**, 036408 (2011).
- [70] Y. Sentoku, K. Mima, S. Kojima, and H. Ruhl, *Phys. Plasmas* **7**, 689 (2000).
- [71] S.V. Bulanov, F. Pegoraro, and A.S. Sakharov, *Phys. Fluids B* **4**, 2499, (1992).
- [72] J. F. Drake, R. G. Kleva, and M. E. Mandt, *Phys. Rev. Lett.* **73**, 1251 (1994).
- [73] D. Biskamp, *Phys. Plasmas* **4**, 1964 (1996).
- [74] D. Biskamp, E. Schwarz, and J. F. Drake, *Phys. Rev. Lett.* **76**, 1264 (1996).
- [75] M.E. Mandt, R.E. Denton, and J.F. Drake, *Geophys. Res. Lett.* **25**, 73 (1998).
- [76] Amita Das and P. H.Diamond, *Phys. Plasmas* **7**, 170 (2000).
- [77] V.S. Lukin, *Phys. Plasmas* **16**, 122105 (2009).
- [78] Ricardo A. Fonseca, et.al., *Phys. Plasmas* **10**, 1975 (2003).
- [79] K.-I. Nishikawa, P. Hardee, G. Richardson, R. Preece, H. Sol, and G.J. Fishman, *ApJ* **622**, 927 (2005).
- [80] A. Hasegawa, and Y. Kodama, *Phys. Rev. Lett.* **41**, 1470 (1978).